

SOME NEW SYNONYMS IN CLYTRINAE
(COLEOPTERA, CHRYSOMELIDAE)

BY F. MONROS

Universidad Nacional de Tucumán, Argentina

I list some new synonyms in Clytrinae, which have been established in the course of my work in connection with Junk's second edition of the "Coleopterorum Catalogus".

Not all the names in this list are plain synonyms; in most cases they represent different color patterns or minor differences in size and microsculpture which tend to be lost in large series. In my opinion, none of these differences is of either specific or subspecific value.

I have seen specimens (in several cases types or specimens from the typical series and typical locality) of all the names listed below.

Macrolenes (Plecomera) quadraticollis Lacord. from South Africa. Lacordaire, 1848, Mon. Phyt. II, p. 105 ♂.

= *M. (P.) varipes* Jacoby, 1901, Trans. Ent. Soc. Lond., 1901, p. 217, pl. 10, f. 5.

Antipus (Gyriodera) cruciata (Thunb.) from South Africa. Thunberg, 1821, Nova Acta Upsala, 8, p. 184.

= *A. (G.) clitellata* Lacordaire, 1848, Mon. Phyt. II, p. 124.

Anomoea laticlavata (Forst.) from the United States and Central America. Forster, 1771, Nov. Spec. Ins., p. 27.

= *A. rufifrons* Lacordaire, 1848, Mon. Phyt. II, p. 135.

= *A. villosa* Jacoby, 1888, Biol. Centr. Amer. Col. 6, 1, Suppl., p. 67, pl. 36, f. 17 ♂.

Diapromorpha (Peploptera) eckloni Lacord. from south east Africa. Lacordaire, 1848, Mon. Phyt. II, p. 244.

= *D. (P.) dorsata* Lacordaire, l.c., p. 245.

Diapromorpha (Peploptera) postica Lacord. from equatorial Africa. Lacordaire, l.c., p. 251.

= *D. (P.) arcuata* Clavareau in Jacoby and Clavareau, 1906, Gen. Ins., fasc. 49, p. 28, footnote.

Diapromorpha (Peploptera) trilineata Lacord. from South Africa. Lacordaire, 1848, Mon. Phyt. II, p. 246.

= *D. (P.) schenklingi* Clavareau, 1909, Ann. Soc. Ent. Belg., 53, p. 386.

- Diapromorpha* (*Peploptera*) *abyssinica* Lef. from Ethiopia.
Lefèvre, 1877, Rev. Mag. Zool., (3) 5, p. 225.
= *D. (P.) schimperi* Lefèvre, 1891, Ann. Soc. Ent. Belg.,
35, c.r. p. CCLI.
- Diapromorpha* (*Peploptera*) *curvilinea* Jac. from South
Africa. Jacoby, 1901, Trans. Ent. Soc. Lond., 1901. p.
222, pl. 10, f. 9.
= *D. (P.) braunsi* Jacoby, 1903, Ent., 36, p. 92.
- Diapromorpha* (*Aspidolopha*) *spilota* (Hope) from Nepal
and Burma. Hope in Gray, 1831, Zool. Misc., p. 30 (*Cryp-
tocephalus*).
= *D. (A.) thoracica* Jacoby, 1892, Ann. Mus. Civ. Geno-
va, 32, p. 879.
- Melitonoma* (*M.*) *decempunctata* (Oliv.), 1808, Ent., 6, pl.
2, f. 24 from tropical and equatorial Africa.
= *M. (M.) patruelis* Lefèvre, 1891, Ann. Soc. Ent. Belg.,
35, c.r. p. CCLII.
= *M. (M.) pedestris* Lefèvre, l.c., p. CCLIII.
= *M. (M.) polysticta* Lacordaire, 1848, Mon. Phyt. II,
p. 372.
= *M. (M.) maculigera* Lacordaire, l.c., p. 376.
= *M. (M.) sobrina* Lacordaire, l.c., p. 377.
= *M. (M.) litigiosa* Lacordaire, l.c., p. 378.
- Megalostomis* (*Megalostomis*) *amazona* Jac. from tropical
America. Jacoby, 1876, Proc. Zool. Soc. Lond., 1876, p.
809.
= *M. (M.) generosa* Baly, 1877, Cist. Ent., 2, p. 181.
= *M. (Minturnia) mariae* Monrós, 1951, Ann. Mag. Nat.
Hist., (12) 4, p. 1154, f. 7-9.
- Megalostomis* (*Megalostomis*) *anachoreta* Lacord. from
Central and northern South America.
Lacordaire, 1848, Mon. Phyt. II, p. 537.
= *M. (Minturnia) chuncho* Monrós, 1951, Ann. Mag.
Nat. Hist., (12) 4, p. 1151, f. 4-5.
= *M. (Minturnia) weyrauchi* Monrós, 1952, Rev. Brasil.
Biol., 12(4), p. 350, f. 2-3.
- Urodera crucifera* Lacord. from the United States to north-
ern Argentina. Lacordaire, 1848, Mon. Phyt. II, p. 454.
= *U. godmani* Jacoby, 1879, Proc. Zool. Soc. Lond., 1879,
p. 775.

Babia quadriguttata (Oliv.) from the United States and Central America. Olivier, 1791, Enc. méth., 6, p. 37.

= *Babia distinguenda* Jacoby, 1888, Biol. Centr. Amer. Col. 6, 1, Suppl., p. 82.

= *Babia amplicollis* Jacoby, l.c., p. 83.

= *Babia parvula* Jacoby, l.c., p. 83.

Babia rufipennis Lacord. from Brazil.

Lacordaire, 1848, Mon. Phyt. II, p. 433.

= *Babia undabunda* Lacordaire, l.c., p. 433.